



RESEARCH ARTICLE

Chemical composition of essential oil from the aerial parts of *Santolina rosmarinifolia* L. a wild Algerian medicinal plant

 Djamel Sarri^{1,*},  Noui Hendel²,  Hadjer Fodil^{1,3},  Giuseppe Ruberto⁴ and  Madani Sarri^{1,5,*}

¹Department of Nature and Life Sciences, Faculty of Sciences, University of M'sila, 28000, M'sila, ALGERIA

²Department of Microbiology and Biochemistry, Faculty of Sciences, University of M'sila, 28000, M'sila, ALGERIA

³Laboratory of Biology, Water and Environment (LBWE), Faculty of Nature and Life Sciences and Earth and Universe Sciences, University of May 8, 1945, 24000, Guelma, ALGERIA

⁴Istituto di Chimica Biomolecolare del Consiglio Nazionale delle Ricerche (ICB-CNR), Via Paolo Gaifami, 18, 95126 Catania, ITALY

⁵Laboratory of Phytotherapy Applied to Chronic Diseases (LPACD), Faculty of Nature and Life Sciences, University of Setif1, 19000, Setif, ALGERIA

*Corresponding authors. Email: djamel.sarri@univ-msila.dz; madani.sarri@univ-msila.dz

Submitted: 21.11.2020; Accepted: 11.01.2021

Abstract

The analysed essential oil in this study was obtained by hydrodistillation from the aerial parts of *Santolina rosmarinifolia* L. (Asteraceae) collected from Hodna area of Algeria. This species is a medicinal herb traditionally used in Algeria. Its essential oil has been analyzed by combining GC-FID and GC-MS. The analysis led to the identification of eighty-two components, representing 91.84% of the whole composition of the sample. The main components were capillene (32.8%), 1,8-cineole (15.1%) and β -myrcene (14.0%).

Keywords: *Santolina rosmarinifolia* L., Asteraceae, Essential oil, GC-MS Analysis, Algeria

Introduction

The *Santolina* L. is a genus belonging to the Asteraceae family (formerly known as the family Compositae, tribe Anthemideae). It is characterized by over ten widely distributed species in the Mediterranean region (Derbesy et al., 1989). Rare in Algeria, the genus *Santolina* is represented by only one species, *Santolina rosmarinifolia* L., called rosemary leaves; a polymorphous plant colonizing forests, pastures and mountainous regions. It is a sub-bushy shrub with closely linear leaves, all tubular flowers, hermaphrodite (peripheral anther sometimes sterile). Inflorescence in dense corymbs, monocle branches and woody stems (Quezel & Santa, 1963). Known in Algeria as Qeissoum and Jaeda (Quezel & Santa, 1963; Baba Aissa, 1999).

Some species of the genus *Santolina* such as *S. chamaecyparissus* L., *S. etrusca* (Lacaita) Marchi & D'Amato, *S. insularis* (Gennari ex Fiori) Arrigoni, *S. neapolitana* Jord. & Fourr. and *S. oblongifolia* Boiss. have been reported in traditional medicine in Spain, Italy, Mediterranean area and India, (Tundis & Loizzo, 2018). In the Algerian traditional pharmacopoeia, *S. rosmarinifolia* L. is used in the treatment of dermatoses in the form of a decoction (Beniston & Beniston, 1984). Also, recommended in the form of flower bouquets arranged in cabinets to protect the linen against moths (Baba Aissa, 1999). These aerial parts are indeed used as vermifuge, stomachic, antispasmodic (Beloued, 2005) and widely pre-registered traditionally as a healing plant (Boudjelal et al., 2013; Sarri et al., 2014). Several *Santolina* species have been studied for their antimicrobial, antifungal, antiviral, and anti-inflammatory activities (Tundis & Loizzo, 2018). The infusion from *S. rosmarinifolia* fresh or dried flower heads was reported to have antipyretic, antihypertensive, hepatoprotective and intestinal anti-inflammatory properties (Ioannou et al., 2007).

This study reports the essential oil composition of *S. rosmarinifolia* from Hammam Dalaa (M'sila Provence) with a distinct chemical profile for the first time.

Materials and Methods

Plant material

The aerial parts of *Santolina rosmarinifolia* L. (Asteraceae) were collected in Hammam Dalaa (M'sila Provence) in May 2017. Avoucher specimen (SD2832/17) was deposited at the Herbarium of the Department of Nature and Life Sciences, Mohamed Boudiaf University (M'sila, Algeria).

Essential oil extraction

Washed aerial parts of the *S. rosmarinifolia* were dried at room temperature in a shady place and then ground to a powder form. The essential oil of the aerial parts of the *S. rosmarinifolia* (300 g) was obtained by hydrodistillation in Clevenger-type apparatus. This operation was carried out for 3 hours. The oil yield was 0.15%, as estimated on the dry weight basis (v/w).

Analysis of the essential oil

On a Hewlett-Packard gas-chromatograph model 5890, fitted with a flame ionization detector, gas chromatographic (GC) analysis was performed (FID). Analyses of GC-FID were carried out under the following analytical conditions: ZB-5 capillary column (30 m x 0.25 mm i.d. x 0.25 μ m film thickness); helium as carrier gas; split mode injection (1:50); 250 and 280°C as injector and detector temperatures, respectively. A programmed oven temperature (40°C to 300 at 2°C/min) was applied.

On the same gas chromatograph, the gas chromatography-mass spectrometry (GC-MS) was performed in connection with a Hewlett-Packard mass spectrometer (model 5971A), 70eV ionization energy, 180°C ion source temperature, mass spectra data were collected over the scan mode in m/z range 40-400.

The individual volatile constituents were identified by comparison of their identical retention indices with those of the compounds known from literature data (Adams, 2007). Furthermore, the chemical identification was confirmed by computer matching of spectral MS data with those stored in the Wiley 275 library and the comparison of the fragmentation patterns with those reported in literature.

Results and Discussion

Chemical composition of the essential oil

Table 1 summarizes the composition and percentage of the essential oil compounds. They have been grouped according to the following classes of components: monoterpene hydrocarbons, oxygenated monoterpenes, sesquiterpenes and others, the last class comprises only the non-terpenoid compounds characterized in the essential oil. On the whole, eighty-two compounds were found, accounting for 91.84% of the total oil.

The most plentiful class in the essential oil of the Algerian *Santolina* was the 'others' amounting to ca. 35% with 17 components, oxygenated monoterpenes was the second class with ca. 22% and 24 components, the following class was monoterpene hydrocarbons with 13 components and an amount of ca. 21%, sesquiterpenes was the last class with the lowest amount (ca. 13%) but with the highest number of components (28). The main constituents were identified as capillene (32.79%), 1,8-cineole (15.08%) and β -myrcene (13.98%).

Species of the genus *Santolina*, particularly *S. rosmarinifolia* and its subspecies, clearly exhibit remarkable variability in terms of essential oil chemical composition (Table 2). This variability is very clear while comparing the major compounds of *S. rosmarinifolia* harvested in a steppe biotope (region of Hodna) which is the subject of this study to that harvested in another forest biotope (Aures region). In this context,

Mehalaine & Chenchouni (2018) showed that certain climatic factors in regions with a semiarid climate had an effect on the quality and composition of the essential oil. This last sample, in fact, showed the presence of two components not found in our sample, namely tricosane (10.6%) and pentacosane (6.7%); a further significant difference was the consistent presence of germacrene-D and α -pinene (30.2 and 10.1%, respectively), together with the absence of capillene (Chibani et al., 2013). A greater similarity with compositional data of our sample has been reported for a sample of *S. rosmarinifolia* ssp. *rosmanifolia* (Palà-Paül et al., 1991, 2001). The phytochemical analysis of *S. rosmarinifolia* collected in Portugal showed that almost all the main compounds found in our sample were present but in lower amount (Ioannou et al., 2007). The flowers oils of *S. rosmarinifolia* collected in Spain presented some similarities when compared with our sample, which in turn did not show the presence of *ar*-curcumene and β -eudesmol (Pérez-Alonso & Velasco-Negueruela, 1988). Finally, some other reports on different species of *Santolina* showed the presence of some components such as 1,8-cineole (Villar et al., 1986; Giner et al., 1933; Cherchi et al., 2001; Flamini & Cioni, 2007; Tirillini et al., 2007; Grosso et al., 2009), β -myrcene (Poli et al., 1997; Zaiter et al., 2015) and capillene (Malti et al., 2019).

Table 1: Chemical composition of essential oil from aerial parts of *Santolina rosmarinifolia* L.

#. ^a	Class/Compounds	RI ^b	% ^c
Monoterpene hydrocarbons			20.9
4	α -Thujene	932	0.1
5	α -Pinene	939	1.3
6	Camphene	955	0.3
7	Thuja-2,4(10)-diene	960	t
9	Sabinene	979	0.7
10	β -Pinene	983	3.5
11	β -Myrcene	997	14.0
12	Phellandrene	1009	t
13	α -Terpinene	1021	0.4
14	<i>p</i> -Cymene	1029	0.3
16	β -Z-Ocimene	1049	t
19	γ -Terpinene	1072	0.3
22	Terpinolene	1093	0.1
Oxygenated monoterpenes			22.3
15	1,8-Cineole	1041	15.1
21	<i>cis</i> -Sabinene hydrate	1089	0.2
23	Perillene	1100	0.5
25	Dehydro sabina ketone	1126	0.1
26	α -Campholenal	1130	0.1
28	<i>trans</i> -Pinocarveol	1146	0.5
29	Camphor	1151	1.0
30	Karahanaenone	1164	t
31	Sabina ketone	1166	t
32	<i>cis</i> -Chrysanthenol	1172	2.0
33	Borneol	1175	0.1
34	<i>cis</i> -Pinocamphone	1182	t
35	Terpinen-4-ol	1186	1.5

36	Thuj-3-en-10-al	1192	0.1
38	α -Terpineol	1199	0.9
39	Myrtenol	1205	0.4
40	<i>trans</i> -Carveol	1227	t
41	<i>cis</i> -Carveol	1237	t
44	Geranial	1278	t
45	<i>p</i> -Menth-1-en-7-al	1283	t
46	<i>cis</i> -Verbenyl acetate	1286	t
47	γ -Terpinen-7-al	1287	t
48	Perilla alcohol	1306	0.1
49	Myrtenyl acetate	1334	t
Sesquiterpenes			13.0
51	α -Copaene	1388	t
52	Bourbonene	1390	t
53	Z-Jasmone	1395	0.1
55	α -Gurjunene	1408	t
56	β -Caryophyllene	1422	t
57	β -Copaene	1431	0.1
58	<i>trans</i> - α -Bergamotene	1445	0.1
59	β -Farnesene	1464	t
60	Sesquisabinene	1471	0.1
61	γ -Gurjunene	1484	0.1
62	γ -Muurolene	1488	0.2
63	Curcumene	1489	0.2
64	Germacrene D	1496	t
66	γ -Cadinene	1526	t
67	Δ -Cadinene	1535	0.4
69	Sphatulenol	1595	3.2
70	Globulol	1598	0.2
71	Salvial-4(14)-en-1-one	1608	0.3
72	Carotol	1618	0.2
73	Oplopenone	1623	0.1
74	β -Atlantol	1626	0.3
75	β -Acorenol	1638	t
76	γ -Eudesmol	1641	t
78	3-Thujopsanone	1657	2.1
79	Eudesma-4(15)-7-dien-1- β -ol	1698	0.3
80	5-neo-Cedranol	1703	0.5
81	Eudesma-4,11-dien-2-ol	1706	3.4
82	β -Acorenone	1708	1.7
Others			35.29
1	3-Hepten-1-ol acetate	773	t
2	Hexanal	803	t
3	<i>E</i> -2-hexenal	859	t
8	Benzaldehyde	966	0.4

17	Benzene acetaldehyde	1058	t
18	2-Octen-2-methyl-6-methylene ^d	1063	0.7
20	Acetophenone	1077	t
24	Isopentylisovalerate	1103	t
27	4- acetyl-1-methylcyclohexane ^d	1135	0.1
37	3,3,5-Trimethyl-1,4-hexadiene ^d	1195	0.1
42	<i>cis</i> -3-Hexenyl isovalerate ^d	1249	0.1
43	<i>p</i> -Ethylacetophenone	1262	t
50	Decanoic acid ^d	1377	t
54	Methyl eugenol	1402	0.1
65	Capillene ^d	1519	32.8
68	Hexenylbenzoate	1580	0.1

^a The numbering refers to elution order on ZB-5 capillary column; ^b Retention index relative to standard mixture of *n*-alkanes on ZB-5 capillary column; ^c Values (area%) represent averages of three determinations (t = trace, <0.05%); ^d Tentatively identified by MS data only.

Table 2. Principal volatile constituents (%) in the essential oils of some specimens of *S. rosmarinifolia* L. and *S. rosmarinifolia* L. ssp. *rosmarinifolia*

Origin / Status / Nature	Part used	Principal components (%)	References
Batna-Aures (Algeria) - <i>Santolina rosmarinifolia</i> L. / Wild	Flowering aerial parts	germacrene-D (30.2), β -myrcene (12.0), tricosane (10.6), β -pinene (10.1), sabinene (7.0) and pentacosane (6.7).	Chibani et al., 2013
Botanical Gardens of Iasi (Romania) / <i>Santolina rosmarinifolia</i> L. / Cultivated	Flower heads	β -eudesmol (13.5), 1,8-cineole (12.9), camphor (8.0), borneol (5.1), <i>ar</i> -curcumene (4.8), terpinen-4-ol (4.5), spathulenol (4.4)	Ioannou et al., 2007
	Leaves	<i>ar</i> -curcumene (9.6), β -phellandrene (8.1), spathulenol (7.5), β -pinene (6.0), γ -muurolene (5.8), myrcene (5.2), camphor (5.2).	
Puente de Madrid (Spain) / <i>Santolina rosmarinifolia</i> L. ssp. <i>rosmarinifolia</i> / Wild	Aerial parts ^(*)	Sabinene (0.3-12), β -pinene (17.0-26.5), myrcene (0.3-15.5), β -phellandrene (14.4-27.6), limonene (3.1-5.0), 1,8-cineole (0.9-1.7), artemisia ketone (1.0-2.4), terpinen-4-ol (0.5-4.0), capillene (t-5.1), <i>ar</i> -curcumene (1.0-2.4), β -eudesmol (0.4-5.0).	Palá-Paúl et al., 2001
Puerto de Nava-cerrad a Madrid (Spain) / <i>Santolina rosmarinifolia</i> L. ssp. <i>rosmarinifolia</i> / Wild	Aerial parts	capillene (35.2), β -phellandrene (14.9), myrcene (13.1), β -pinene (7.8), sabinene (5.5), <i>ar</i> -curcumene (4.3)	Palá-Paúl et al., 1999
El Escorial and Puerto de Galapagar a Madrid (Spain) / <i>Santolina rosmarinifolia</i> L. / Wild	Flowers	1,8-cineole (8.9), β -pinene (8.9), myrcene (7.8), sabinene (6.6), <i>ar</i> -curcumene (5.8), β -eudesmol (13.4).	Pérez-Alonso & Velasco-Negueruela, 1988

(*) Components of the EO (in %) of *Santolina rosmarinifolia* L. ssp. *rosmarinifolia* over one year (min-max); t=traces

Conclusion

This study forms the first report on the chemical composition of the essential oil from *Santolina rosmarinifolia* L. harvested in a steppe biotope. The obtained results could contribute to the valorisation of this Algerian medicinal plant.

ACKNOWLEDGMENT

The authors are grateful to CNEPRU (D01N01UN280120150001) and we wish to thank Consiglio Nazionale delle Ricerche (C.N.R., Rome) for financial support. The authors would also to thank Mr. A. MAIZA and N. BEZAI for the technical support.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

REFERENCES

- Adams, R.P. (2001). *Identification of essential oil components by gas chromatography/mass spectroscopy*. Illinois, USA: Allured publishing Co. Carol Stream.
- Baba Aissa F., 1999. *Encyclopédie des plantes utiles : flore d'Algérie et du Maghreb*, Librairie moderne, Rouiba, Algérie, 368p.
- Beloued, A. (2005). *Plantes médicinales d'Algérie*. Alger, Algérie: Office des Publications Universitaire (OPU).
- Beniston, N.T. & Beniston, W.S. (1984). *Fleurs d'Algérie*. Alger, Algérie: Entreprise Nationale du Livre (ENL).
- Boudjelal, A., Henchiri, C., Sari, M., Sarri, D., Hendel, N., Benkhaled, A., Ruberto, G. (2013). Herbalists and wild medicinal plants in M'Sila (North Algeria): An ethnopharmacology survey. *Journal of Ethnopharmacology*, 148(2), 395-402.
- Cherchi, G., Deidda, D., De Gioannis, B., Marongiu, B., Pompei, R., Porcedda, S. (2001). Extraction of *Santolina insularis* essential oil by supercritical carbon dioxide: influence of some process parameters and biological activity. *Flavour and Fragrance Journal*, 16(1), 35-43.
- Chibani, S., Labed, A., Kabouche, A., Semra, Z., Smati, F., Aburjai, T., Kabouche, Z. (2013). Antibacterial activity and chemical composition of essential oil of *Santolina rosmarinifolia* L. (Asteraceae) from Algeria. *Der Pharmacia Letter*, 5(2), 238-241.
- Derbesy, M., Touche, J., Zola, A. (1989). The essential oil of *Santolina chamaecyparissus* L. *Journal of Essential Oil Research*, 1(6), 269-275.
- Flamini, G., Cioni, P.L. (2007). Seasonal variation of the chemical constituents of the essential oil of *Santolina etrusca* from Italy. *Chemistry and Biodiversity*, 4(5), 1008-1019.
- Giner, R.M., Manez, S., Rios, J.L. (1993). Seasonal variations in the essential oil of *Santolina chamaecyparissus* L. *Scientia Pharmaceutica*, 61, 169-173.
- Grosso, C., Figueiredo, A.C., Burillo, J., Mainar, A.M., Urieta, J.S., Barosso, J.G., Coelho, J.A., Palavra, A.M.F. (2009). Enrichment of the thymoquinone content in volatile oil from *Satureja montana* using supercritical fluid extraction. *Journal of Separation Science*, 32, 328-334.
- Ioannou, E., Poiata, A., Hancianu, M., Tzakou, O. (2007). Chemical composition and *in vitro* antimicrobial activity of the essential oils of flower heads and leaves of *Santolina rosmarinifolia* L. from Romania. *Natural Product Research*, 21(1), 18-23.
- Malti, C.E.W., Baccati, C., Mariani, M., Hassani, F., Babali, B., Atik-Bekkara, F., Paoli, M., Maury, J., Tomi, F., Bekhechi, C. (2019). Biological activities and chemical composition of *Santolin africana* Jord. et Fourr. aerial part essential oil from Algeria: occurrence of polyacetylene derivatives. *Molecules*, 24(1), 204.
- Mehalaine, S. & Chenchouni, H. (2018). Effect of climatic factors on essential oil accumulation in two Lamiaceae species from Algerian semiarid lands. *Conference of the Arabian Journal of Geosciences*, 57-60.
- Palá-Paúl, J., Pérez-Alonso, M.J., Velasco-Negueruela, A., Pala-Paul, R., Sanz Fco Conejero, J. (2001). Seasonal variation in chemical constituents of *Santolina rosmarinifolia* L. ssp. *Rosmarinifolia*. *Biochemical Systematics and Ecology*, 29(7), 663-672.
- Palá-Paúl, J., Pérez-Alonso, M.J., Velasco-Negueruela, A., Ramos-Vázquez, P., Gómez-Contreras, F., Sanz J. (1999). Essential oil of *Santolina rosmarinifolia* L. ssp. *rosmarinifolia*: first isolation of capillene, a diacetylene derivative. *Flavour and Fragrance Journal*, 14(2), 131-134.

- Pérez-Alonso, M.J., Velasco-Negueruela, A. (1988). The essential oils of four *Santolina* species. *Flavour and Fragrance Journal*, 3(1), 37-42.
- Poli, F., Bonsignore, L., Loy G., Sacchetti, G., Ballero, M. (1997). Comparison between the essential oils of *Santolina insularis* (Genn. ex Fiori) Arrigoni and *Santolina corsica* Jord. et Fourr. from the island of Sardinia (Italy). *Journal Ethnopharmacology*, 56(3), 201-208.
- Quézel, P. & Santa, S. (1963). *Nouvelle flore de l'Algérie et des régions désertiques méridionales*. Paris, France : Centre National de la Recherche Scientifique (CNRS).
- Sarri, M., Mouyet, F.Z., Benziane, M., Cheriet, A. (2014). Traditional use of medicinal plants in a city at steppic character (M'sila, Algeria). *Journal of Pharmacy and Pharmacognosy Research*, 2(2), 31-35.
- Tirillini, B., Ricci, A., Pintore, G., Chessa, M., Menghini, L., Pagiotti, R. (2007). Essential oil composition of *Santolina etrusca* from Italy. *Chemistry of Natural Compounds*, 43(1), 44-46.
- Tundis, R. & Loizzo, M.R. (2018). A review of the traditional uses, phytochemistry and biological activities of the genus *Santolina*. *Planta Medica*, 84, 627-637
- Villar, A., Giner, R.M., Rios, J.L. (1986). Chemical composition of *Santolina chamaecyparissus* ssp. *squarrosa* essential oil. *Journal of Natural Products*, 49(6), 1143-1144.
- Zaiter, L., Benayache, F., Beghidja, N., Figueredo, G., Chalard, P., Chalchat, J.C. (2015). Essential oils of *Santolina africana* Jord. & Fourr. and *Santolina chamaecyparissus* L. *Journal of Essential Oil Bearing Plants*, 18(6), 1338-1342.